


IN THE CLAIMS

Please amend claim 1 as follows:

Claim 1. (Currently Amended). An active-matrix display containing a ferroelectric (chiral smetic) liquid-crystal mixture in a liquid crystal layer in the form of a monodomain having an unambiguously defined direction of the layer normal z of the SmC^* phase, where the layer normal z and the preferential direction n of the nematic or cholesteric phase (N^* phase) form an angel of more than 5° , wherein the liquid-crystal mixture ~~comprising~~ comprises at least one compound of the formula (1)



where the symbols are as defined below:

R^1 , R^2 are, independently of one another, identical or different and are each

a) hydrogen, fluorine or CN

a straight-chain or branched alkenyl, alkenyloxy, alkyl or alkyloxy radical (with or without asymmetric carbon atoms) having 2 to 16 carbon atoms, where

b1) one or two nonterminal $\text{-CH}_2\text{-}$ groups may be replaced by -O- , -OC(=O)- , -C(=O)- , -C(=O)O- , $\text{-Si(CH}_3\text{)}_2\text{-}$, -CH(CH)- and/or one or two $\text{-CH}_2\text{-}$ groups may be replaced by -CH=CH- or $\text{-C}\equiv\text{C-}$

and one or more H atoms may be replaced by F and/or

b2) one or more $\text{-CH}_2\text{-}$ groups may be replaced by phenylene-1,4-diyl (unsubstituted, monosubstituted or disubstituted by F), phenylene-1,3-diyl (unsubstituted, monosubstituted or disubstituted by F), cyclohexane-

1,4-diyl (unsubstituted or monosubstituted by F or CN) or cyclopropane-

1,2-diyl

and one or more H atoms may be replaced by F

with the provisos that only one of the radicals R^1 , R^2 can be hydrogen, F or CN and that two adjacent $-CH_2-$ groups cannot be replaced by $-O-$

M^1 , M^2 are, independently of one another, identical or different and are each

$-C(=O)O-$, $-OC(=O)-$, $-CH_2O-$, $-OCH_2-$, $-CF_2O-$, $-OCF_2-$, $-CH_2CH_2-$, $-CF_2CF_2-$,
 $-CH=CH-$, $-CH=CF-$, $-CF=CF-$, $-C\equiv C-$, $-CH_2CH_2C(=O)O-$, $-OC(=O)CH_2CH_2-$,
 $-(CH_2)_4-$, $-OCH_2CH_2CH_2-$, $-CH_2CH_2CH_2O-$, $-OCH_2CF_2CH_2-$, $-CH_2CF_2CH_2O-$ or a
single bond

A^1 , A^2 , A^3 are, independently of one another, identical or different and are each

cyclohexane-1,4-diyl (unsubstituted or monosubstituted by F, CH_3 , CN),
cyclohex-1-ene-1,4-diyl, cyclohex-2-ene-1,4-diyl, 2-oxocyclohexane-1,4-diyl,
2-cyclohexen-1-one-3,6-diyl, 1-alkyl-1-silacyclohexane-1,4-diyl, bicyclo[2.2.2]-
octane-1,4-diyl, spiro[4.5]decane-2,8-diyl, spiro[5.5]undecane-3,9-diyl,
phenylene-1,4-diyl (unsubstituted, monosubstituted or disubstituted by CN, CH_3 ,
 CF_3 , OCH_3 , unsubstituted, monosubstituted, disubstituted, trisubstituted or
tetrasubstituted by F), phenylene-1,3-diyl (unsubstituted, monosubstituted or
disubstituted by CN, CH_3 , CF_3 , OCF_3 , unsubstituted, monosubstituted,
disubstituted, trisubstituted or tetrasubstituted by F), thiophene-2,5-diyl,
thiophene-2,4-diyl, (1,3,4)-oxadiazole-2,5-diyl, (1,3,4)-thiadiazole-2,5-diyl,

1,3-thiazole-2,5-diyl, 1,3-thiazole-2,4-diyl, (1,3)-oxazole-2,5-diyl, isoxazole-2,5-diyl, indane-2,6-diyl, naphthalene-2,6-diyl (unsubstituted, monosubstituted or disubstituted by F or CN), 1,2,3,4-tetrahydronaphthalene-2,6-diyl, decaline-2,6-diyl, pyrimidine-2,5-diyl (unsubstituted or monosubstituted by F), pyridine-2,5-diyl (unsubstituted, monosubstituted or disubstituted by F), pyrazine-2,5-diyl (unsubstituted or monosubstituted by F), pyridazine-3,6-diyl, quinoline-2,6-diyl, quinoline-3,7-diyl, isoquinoline-3,7-diyl, quinazoline-2,6-diyl, 5,6,7,8-tetrahydroquinazoline-2,6-diyl, quinoxaline-2,6-diyl, 1,3-dioxane-2,5-diyl (unsubstituted or monosubstituted by CN), benzothiazole-2,6-diyl, piperidine-2,4-diyl, piperazine-1,4-diyl

B¹ is cyclohexane-1,4-diyl (unsubstituted, monosubstituted or disubstituted by F, CH₃, CN), perfluorocyclohexane-1,4-diyl, cyclohex-1-ene-1,4-diyl, cyclohex-2-ene-1,4-diyl, 1-alkyl-1-silacyclohexane-1,4-diyl, bicyclo[2,2,2]octane-1,4-diyl, cyclopentane-1,3-diyl, cycloheptane-1,4-diyl, tetrahydrofuran-2,5-diyl, tetrahydrofuran-2,4-diyl, phenylene-1,4-diyl (unsubstituted, monosubstituted or disubstituted by CN, CH₃, CF₃, OCF₃, unsubstituted, monosubstituted, disubstituted, trisubstituted or tetrasubstituted by F), phenylene-1,3-diyl (unsubstituted, monosubstituted or disubstituted by CN, CH₃, CF₃, OCF₃, unsubstituted, monosubstituted, disubstituted or trisubstituted by F), thiophene-2,5-diyl (unsubstituted or monosubstituted by F), thiophene-2,4-diyl (unsubstituted or monosubstituted by F), 1,3-thiazol-2,5-diyl (unsubstituted or monosubstituted by F), 1,3-thiazol-2,4-diyl (unsubstituted or monosubstituted by F), (1,3,4)-thiadiazol-

2,5-diyl, 1,3-dioxane-2,5-diyl (unsubstituted or monosubstituted by CN), tetrahydro-
pyran-2,5-diyl, 6,6-difluorotetrahydro pyran-2,5-diyl, 6,6-difluoro-
2,3-dihydro-6H-pyran-2,5-diyl, 6-fluoro-3,4-dihydro-2H-pyran-2,5-diyl,
piperidine-1,4-diyl, piperazine-1,4-diyl, pyrimidine-2,5-diyl (unsubstituted or
monosubstituted by F), pyridine-2,5-diyl (unsubstituted or monosubstituted by F),
1,2,3,4-tetrahydronaphthalene-2,6-diyl, decaline-2,6-diyl

B² is cyclohexane-1,4-diyl (unsubstituted, monosubstituted or disubstituted by F,
CH₃, CN), cyclohex-1-ene-1,4-diyl (unsubstituted or monosubstituted by F),
cyclohex-2-ene-1,4-diyl, 1-alkyl-1-silacyclohexane-1,4-diyl, bicyclo[2.2.2]-
octane-1,4-diyl, phenylene-1,4-diyl (unsubstituted, monosubstituted or
disubstituted by CN, CH₃, CF₃, OCF₃, unsubstituted, monosubstituted,
disubstituted, trisubstituted or tetrasubstituted by F), phenylene-1,3-diyl
(unsubstituted, monosubstituted or disubstituted by CN, CH₃, CF₃, OCF₃,
unsubstituted, monosubstituted, disubstituted or trisubstituted by F), thiophene-
2,5-diyl, thiophene-2,4-diyl, 1,3-thiazole-2,5-diyl, 1,3-thiazole-2,4-diyl, (1,3,4)-
thiadiazole-2,5-diyl, 1,3-dioxane-2,5-diyl (unsubstituted or monosubstituted by
CN), tetrahydrofuran-2,5-diyl, tetrahydropyran-2,5-diyl, 6,6-difluoro-
tetrahydropyran-2,5-diyl, 6,6-difluoro-2,3-dihydro-6H-pyran-2,5-diyl, 6-fluoro-
3,4-dihydro-2H-pyran-2,5-diyl, pyrimidine-2,5-diyl (unsubstituted or mono-
substituted F), pyridine-2,5-diyl (unsubstituted or monosubstituted F), indane-
2,6-diyl, piperidine-1,4-diyl, piperazine-1,4-diyl, pyrimidine-2,5-diyl
(unsubstituted or monosubstituted by F)

X is $-(CH_2)_n-$, where

- a) one or two $-CH_2-$ groups may be replaced by $-O-$ or $-C(=O)-$ and/or
- b) one $-CH_2CH_2-$ group may be replaced by $-CH=CH-$ and one or more H of the $-CH_2-$ groups may be replaced by F

with the provisos that

- 1) n is 2, 3 or 4
- 2) two adjacent $-CH_2-$ groups cannot be replaced by $-O-$

a, b, c are each zero, 1 or 2, with the provisos that

- 1) a must be 1 when R^1 is hydrogen, F or CN
- 2) the sum of a+b+c is at least 1
- 3) the radicals A and M, respectively, in the brackets may be identical or different when the corresponding index is 2 in a liquid-crystal layer in the form of a monodomain having an unambiguously defined direction of the layer normal z of the SmC* phase, where the layer normal z and the preferential direction n of the nematic or cholesteric phase (N* phase) form an angle of more than 5°.

Claim 2. (Cancelled). An active-matrix display as claimed in claim 1, containing a liquid-crystal layer in the form of a monodomain having an unambiguously defined direction of the layer normal z of the SmC* phase, where the layer normal z and the preferential direction n of the nematic or cholesteric phase (N* phase) form an angle of more than 5°, and the liquid-crystal

layer is composed of a ferroelectric (chiral smectic) liquid-crystal mixture comprising at least one compound of the formula (I).

Claim 3. (Previously amended). A display as claimed in claim 1, wherein the liquid-crystal mixture has a spontaneous polarization of $< 200 \text{ nC/cm}^2$ and $DT(15,1) > 20$.

Claim 4. (Previously amended). A display as claimed in claim 1, wherein, in (I),

X is $-\text{OC}(=\text{O})-$, $-\text{OCH}_2-$ or $-\text{OC}(=\text{O})\text{CH}_2\text{CH}_2-$.

Claim 5. (Previously Amended). A display as claimed in claim 1, wherein, in (I),



B^1 is cyclohexane-1,4-diyl, cyclohex-1-ene-1,4-diyl, phenylene-1,4-diyl,
unsubstituted, monosubstituted or disubstituted by F, or thiophene-2,5-diyl.

Claim 6. (Previously Amended). A display as claimed in claim 1, wherein, in (I),

A^1 is pyrimidine-2,5-diyl (unsubstituted or monosubstituted by F), pyridine-2,5-diyl
(unsubstituted or monosubstituted by F), phenylene-1,4-diyl (unsubstituted,
monosubstituted or disubstituted by F), or (1,3,4)-thiadiazol-2,5-diyl.








Please amend claim 7 as follows:


Claim 7. (Previously Amended). A display as claimed in claim 1, wherein the liquid-crystal mixture is composed of 3 to 30 compounds and comprises at least one compound of the formula (I) and at least one compound of the formula (II) below and, if desired, at least one compound of the formula (III) below



where:

R^{10} , R^{11} are as defined for R^1 , R^2 , where additionally the terminal $-CH_3-$ group may in each case be replaced by one of the chiral groups (optically active or racemic) below:





R^3, R^4, R^5, R^6, R^7 are identical or different and are each

- a) hydrogen
- b) a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 16 carbon atoms, where
 - b1) one or more nonadjacent and nonterminal CH_2 groups may be replaced by $-O-$ and/or
 - b2) one or two CH_2 groups may be replaced by $-CH=CH-$.

- c) R^4 and R^5 together may alternatively be $-(CH_2)_4-$ or $-(CH_2)_5-$ if they are attached to an oxirane, dioxolane, tetrahydrofuran, tetrahydropyran, butyrolactone or valerolactone system;

R^{12} is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 16 carbon atoms, where one or more H may be replaced by F and one or two nonadjacent nonterminal $-CH_2-$ groups may be replaced by $-O-$

$Z^1, Z^2, Z^3, Z^4, Z^5, Z^6$ are each, independently of one another, H or F



is a bivalent radical selected from the group consisting of pyridine-2,5-diyl, unsubstituted or monosubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, pyrazine-2,5-diyl, unsubstituted or monosubstituted by F,







is a bivalent radical selected from the group consisting of cyclohexane-1,4-diyl, unsubstituted or monosubstituted by CN, CH_3 , or disubstituted by F, cyclohex-1-ene-1,4-diyl, perfluorocyclohexane-1,4-diyl, cyclohex-2-ene-1,4-diyl, 1-alkyl-1-silacyclohexane-1,4-diyl, bicyclo[2.2.2]octane-1,4-diyl.

Please amend claim 8 as follows:

Claim 8. (Currently Amended). A display as claimed in claim 4 7, wherein the liquid-crystal mixture is composed of 3 to 30 compounds and comprises at least one compound of the


formula (I) and at least one compound of the formula (II) and at least one additional compound, selected from the group consisting of (III), (IV), (V), (VI), (VII), where the compounds of the formulae (II) and (III) are as defined in claim 7,




where the symbols and indices are as defined in claim 7.

Claim 9. (Previously Amended). A display as claimed in claim 1, wherein the liquid-crystal mixture is composed of 3 to 30 compounds and comprises at least one compound of the formula (I) and at least one compound of the formula (II) and at least one additional compound








selected from the group consisting of (VIII), (IX), (X), (XI), (XII), (XIII), (XIV), (XV), (XVI), (XVII), where the compounds of the formulae (II) and (III) are as defined in claim 7.




(VIII)




(IX)





(XVII)


where the symbols and indices are as defined in claim 7 or as defined below:



is a bivalent radical selected from the group consisting of naphthalene-2,6-diyl, in which one or two ring carbon atoms may be replaced by N and which can be monosubstituted or disubstituted by F or CN and in which D¹ or D² may also be a (saturated) alicycle



is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by CN, or unsubstituted, monosubstituted, disubstituted, trisubstituted or tetrasubstituted by F, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, cyclohexane-1,4-diyl




is a bivalent radical selected from the group consisting of indane-2,5-diyl, unsubstituted, monosubstituted or disubstituted by F in the aromatic ring, indan-1-one-2,6-diyl, unsubstituted, monosubstituted or disubstituted by F in the aromatic ring,


benzothiazole-2,6-diyl, benzothiazole-2,5-diyl, benzo[b]thiophene-2,5-diyl, benzo[b]-
thiophene-2,6-diyl




is a bivalent radical selected from the group consisting of (1,3,4)-
thiadiazole-2,5-diyl, (1,3)-thiazole-2,5-diyl, thiophene-2,5-diyl, (1,3,4)-oxadiazole-
2,5-diyl, (1,3)-oxazole-2,5-diyl, isoxazole-2,5-diyl



is a bivalent radical selected from the group consisting of 1,1'-
biphenyl-4,4'-diyl, unsubstituted, monosubstituted or disubstituted by C^N, or
unsubstituted, monosubstituted, disubstituted, trisubstituted or tetrasubstituted by F,
1,1'-phenylcyclohexyl-4,4'-diyl, 5,5'-pyridylpyrimidine-2,2'-diyl, unsubstituted or
monosubstituted by F in one or both of the heterocycles, 5,2'-pyridylpyrimidine-
2,5'-diyl, unsubstituted or monosubstituted by F in one or both of the heterocycles,
1,2'-phenyldioxane-4,5'-diyl, 1,2'-(2-fluorophenyl)dioxane-4,5'-diyl,
1,2'-(3-fluorophenyl)dioxane-4,5'-diyl, 1,2'-(2,3-difluorophenyl)dioxane-4,5'-diyl



is a bivalent phenanthrene-2,7-diyl radical in which one or two
ring carbon atoms may be replaced by N and which may be monosubstituted,
disubstituted, trisubstituted or tetrasubstituted by F and in which P² and/or P³ may be a
(saturated) alicycle



is a bivalent fluorene-2,7-diyl radical in which the $-\text{CH}_2-$ group in U^2 may be replaced by $-\text{C}(=\text{O})-$, $-\text{C}(\text{H})-$ or $-\text{C}(\text{F})_2-$.



is a bivalent radical selected from the group consisting of phenylene-1,3-diyl, unsubstituted, monosubstituted or disubstituted by F, cyclohexane-1,3-diyl, unsubstituted or monosubstituted by F or CN, pyridine-2,6-diyl, pyridine-2,4-diyl, pyridine-3,5-diyl, pyridine-4,6-diyl, pyrimidine-4,6-diyl.



is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by CN or F, naphthalene-2,6-diyl, in which one or two ring carbon atoms may be replaced by N and which may be monosubstituted or disubstituted by CN or F, cyclohexane-1,4-diyl, cyclohex-1-ene-1,4-diyl, bicyclo[2.2.2]octane-1,4-diyl, (1,3)-dioxane-2,5-diyl, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, (1,3,4)-thiadiazole-2,5-diyl, indane-2,5-diyl, unsubstituted, monosubstituted or disubstituted by F in the aromatic ring, thiophene-2,5-diyl



is a bivalent radical selected from the group consisting of phenylene-1,4-diyl, unsubstituted, monosubstituted or disubstituted by CN or F, naphthalene-2,6-diyl, in which one or two ring carbon atoms may be replaced by N and which may be

monosubstituted or disubstituted by CN or F, cyclohexane-1,4-diyl, cyclohex-1-ene-1,4-diyl, bicyclo[2.2.2]octane-1,4-diyl, (1,3)-dioxane-2,5-diyl, indane-2,5-diyl, unsubstituted, monosubstituted or disubstituted by F in the aromatic ring, thiophene-2,5-diyl

p, q, s are each zero or 1

r is 1 or 2.


Please amend claim 10 as follows:

Claim 10. (Currently Amended). ~~A~~ The display as claimed in claim 1, wherein the smectic liquid-crystal mixture ~~as claimed in claim 1 comprising~~ comprises from 10 to 60% of one or more compounds of the formula (I).

Please amend claim 11 as follows:

Claim 11. (Currently Amended). ~~A~~ The display as claimed in claim 1 wherein the chiral smectic liquid-crystal mixture ~~as claimed in claim 1 comprising~~ comprises from 10 to 60% of 1 to 15 compounds of the formula (I) and from 40 to 90% of 2 to 15 compounds of the formula (II).

Claim 12. (Previously Amended). A compound selected from compounds of the formula (XXIII), where:



n	9	9	9	9	9	9	9	9	10	10	10	10	10	10	10	11	11	11	11	11	11				
m	3	4	5	6	7	8	9	10	11	3	4	6	7	8	9	10	11	12	5	6	7	8	9	10	11
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-


n	12	12	12	12	12	12	13	13	13	13	13	13	13	13	14	14	14	14	14	14					
m	4	6	7	8	9	10	11	3	4	5	6	7	8	9	10	11	3	4	5	6	7	8			
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

n	14	14	14	14	6	6	6	6	6	6	6	6	6	6	7	7	7	7	7	7	7	7	7	8	8	
m	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4
X	-	-	-	-	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	8	8	8	8	8	8	8	9	9	9	9	9	9	9	9	9	10	10	10	10	10	10	10	10	10	
m	5	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	11	11	11	11	11	11	11	11	11	12	12	12	12	12	12	12	12	12	13	13	13	13	13	13	13	
m	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

compounds of the formula (XXIV), where:




n is an integer from 8 to 14

m is an integer from 3 to 11

X is a single bond

with the exception of n=11, m=3 or 5, X is a single bond,

compounds of the formula (XXV), where:




n is an integer from 2 to 13

m is an integer from 3 to 11

X is O or a single bond

with the exception of n=2, m=11, X=O; n=5, m=5, X=O,

compounds of the formula (XXVI), where:




n is an integer from 5 to 13

m is an integer from 3 to 10

with the exception of n=8, m=5,

compounds of the formula (XXVII), where:



n	8	8	8	8	8	8	8	9	9	9	9	9	9	9	9	10	10	10	10	10	10	10	10	
m	3	4	6	7	8	9	10	3	4	5	6	7	8	9	10	11	3	4	5	6	7	8	9	10
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-


n	11	11	11	11	11	11	11	11	12	12	12	12	12	12	12	12	13	13	13	13	13	13	13
m	3	4	5	6	7	8	9	10	3	4	5	6	7	8	9	10	11	3	4	5	6	7	8
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

n	13	13	13	14	14	14	14	14	14	14	7	7	7	7	7	7	7	7	8	8	8	8	8	
m	9	10	11	3	4	5	6	7	8	9	10	3	4	5	6	7	8	9	10	11	3	4	6	7
X	-	-	-	-	-	-	-	-	-	-	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	8	8	8	8	9	9	9	9	9	9	9	9	9	10	10	10	10	10	10	10	11	11	11	
m	8	9	10	11	3	4	5	6	7	8	9	10	11	3	4	5	6	7	8	9	10	3	4	5
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	11	11	11	11	11	11	11	12	12	12	12	12	12	12	12	13	13	13	13	13	13	13	13	
m	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	3	4	5	6	7	8	9	10	11
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

compounds of the formula (XXIX), where:




n	6	6	6	7	7	7	7	7
m	7	8	9	4	6	8	9	10
X	-	-	-	-	-	-	-	-

n	8	8	9	9	9	9	9	9	9	10	10	10
m	8	10	3	4	6	7	8	9	10	8	9	19
X	-	-	-	-	-	-	-	-	-	-	-	-

n	6	6	6	6	6	6	6	7	7	7	7	7	7	7	
m	3	4	6	7	8	9	10	3	4	5	6	7	8	9	10
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	8	8	8	8	8	8	8	8	9	9	9	9	9	9	10	10	10	10	10	10	10
m	3	4	5	6	7	8	9	10	5	6	7	8	9	10	4	5	6	7	8	9	10
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

compounds of the formula (XXX), where:




n is an integer from 5 to 13

m is an integer from 3 to 10

with the exception of n=8, m=4; n=9, m=3.

Claim 13. (Previously Amended). A compound selected from compounds of the formula (XXX1), where:



n	10	10	10	10	10	10	10	10	10	10	11	11	11	11	11	11	11	11	12	12	12	12		
m	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

n	12	12	12	12	12	12	13	14	14	14	14	14	14	14	14	14	6	6	6	6	6	6	6	6
m	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	O	O	O	O	O	O	O	O


n	6	6	7	7	7	7	7	7	7	7	7	8	8	8	8	8	8	8	8	8	9	9	9	9
m	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	8	9	10	11	12	3	4	5
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	9	9	9	9	9	9	9	10	10	10	10	10	10	10	10	10	10	11	11	11	11	11	11	11
m	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	11	11	11	12	12	12	12	12	12	12	12	12	13	13	13	13	13	13	13	13	13	14		
m	10	11	12	3	4	5	6	7	8	9	10	11	12	3	4	5	6	7	8	9	10	11	12	3
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	14	14	14	14	14	14	14	14	14
m	4	5	6	7	8	9	10	11	12
X	O	O	O	O	O	O	O	O	O

compounds of the formula (XXVIII), where:




n	11	12	13	14	13	14	12	13	14	13	14	10	11	12	13	14	13	14	9	10	11	12	13	10
m	5	5	5	5	6	6	7	7	7	8	8	9	9	9	9	9	10	10	11	11	11	11	11	12
X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

n	11	12	13	14	6	6	6	6	6	6	6	6	7	7	7	7	7	7	7	7	8	8		
m	12	12	12	12	4	5	6	7	8	9	10	11	12	4	5	6	7	8	9	10	11	12	4	6
X	-	-	-	-	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	8	8	8	8	9	9	9	9	9	9	9	10	10	10	10	10	10	10	10	10	11	11	11	11
m	8	10	11	12	4	5	6	8	9	10	11	12	4	5	7	8	9	10	11	12	4	5	6	7
X	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O	O

n	11	11	11	11	11	12	12	12	12	12	12	12	12
m	8	9	10	11	12	5	6	7	8	9	10	11	12
X	O	O	O	O	O	O	O	O	O	O	O	O	O

compounds of the formula (XXXII), where:



n	5	5	5	5	5	5	5	5	6	6	6	6	6	6	6	7	7	7	7	
m	2	3	4	5	6	7	8	9	2	3	4	5	6	7	8	9	2	3	4	5

n	7	7	7	7	8	8	8	8	8	8	8	9	9	9	9	9	9	9	10	10	10	10			
m	6	7	8	9	2	3	4	5	6	7	8	9	2	3	4	5	6	7	8	9	2	3	4	5	6

n	10	10	10	11	11	11	11	11	11	11	11	12	12	12	12	12	12	12	13	13	13	
m	7	8	9	2	3	4	5	6	7	8	9	2	3	4	5	6	7	8	9	2	3	4

n	13	13	13	13	13	14	14	14	14	14	14	14	14
m	5	6	7	8	9	2	3	4	5	6	7	8	9

and where Z is H or F in all cases.